



# The Brazilian Journal of INFECTIOUS DISEASES

[www.elsevier.com/locate/bjid](http://www.elsevier.com/locate/bjid)



## Original article

# Antifungal activities of tacrolimus in combination with antifungal agents against fluconazole-susceptible and fluconazole-resistant *Trichosporon asahii* isolates



Thaís Felli Kubiça<sup>a,b</sup>, Laura Bedin Denardi<sup>a</sup>, Maria Isabel Azevedo<sup>c</sup>, Vanessa Oliveira<sup>c</sup>,  
Luiz Carlos Severo<sup>e</sup>, Janio Moraes Santurio<sup>c,d</sup>, Sydney Hartz Alves<sup>a,c,d,\*</sup>

<sup>a</sup> Universidade Federal de Santa Maria (UFSM), Centro de Ciências da Saúde, Programa de Pós-Graduação em Ciências Farmacêuticas, Santa Maria, RS, Brazil

<sup>b</sup> Universidade Regional Integrada do Alto Uruguai e das Missões (URI), Santiago, RS, Brazil

<sup>c</sup> Universidade Federal de Santa Maria (UFSM), Centro de Ciências da Saúde, Programa de Pós-Graduação em Farmacologia, Santa Maria, RS, Brazil

<sup>d</sup> Universidade Federal de Santa Maria (UFSM), Departamento de Microbiologia e Parasitologia, Santa Maria, RS, Brazil

<sup>e</sup> Complexo Hospitalar Santa Casa, Porto Alegre, RS, Brazil

## ARTICLE INFO

### Article history:

Received 14 March 2016

Accepted 5 August 2016

Available online 30 September 2016

### Keywords:

Susceptibility

FK506

Resistance

Synergism

## ABSTRACT

The antifungal activity of tacrolimus in combination with antifungal agents against different fungal species has been previously reported. Here we report the *in vitro* interactions between tacrolimus and amphotericin B, fluconazole, itraconazole, and caspofungin against 30 clinical isolates of both fluconazole-susceptible and fluconazole-resistant *Trichosporon asahii*. For these analyses, we used the broth microdilution method based on the M27-A3 technique and checkerboard microdilution method. Tacrolimus showed no activity against *T. asahii* strains (minimal inhibitory concentrations, MICs > 64.0 µg mL<sup>-1</sup>). However, a larger synergistic interaction was observed by the combinations tacrolimus + amphotericin B (96.67%) and tacrolimus + caspofungin (73.33%) against fluconazole-susceptible isolates. Combinations with azole antifungal agents resulted in low rates of synergism for this group (fluconazole + tacrolimus = 40% and itraconazole + tacrolimus = 10%). Antagonistic interactions were not observed. For the fluconazole-resistant *T. asahii* group, all tested combinations showed indifferent interactions. The synergism showed against fluconazole-susceptible *T. asahii* isolates suggests that the potential antifungal activity of tacrolimus deserves *in vivo* experimental investigation, notably, the combination of tacrolimus with amphotericin B or caspofungin.

© 2016 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [sydneyalves.ufsm@gmail.com](mailto:sydneyalves.ufsm@gmail.com) (S.H. Alves).

<http://dx.doi.org/10.1016/j.bjid.2016.08.008>

1413-8670/© 2016 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The incidence of invasive mycoses caused by emergent fungal pathogens has proportionally grown with the increased number of immunocompromised hosts, such as AIDS patients, transplant recipients treated with immunosuppressive drugs, and those on cancer therapy.<sup>1-3</sup> *Trichosporon asahii* is the most frequently involved species in disseminated and deep-seated trichosporonosis including systemic infections due to therapeutic failure in transplanted patients.<sup>2,3</sup>

The first-line treatment for trichosporonosis includes the use of azole antifungal agents, since *Trichosporon* spp. is resistant to amphotericin B and echinocandins.<sup>4-6</sup> However, the frequent exposure to azoles can lead to development of secondary resistance to azoles and sometimes to multidrug resistance, resulting in therapeutic failures<sup>7-9</sup> and increasing mortality rates.<sup>1,10</sup> Combination therapy is a rational alternative that has been studied to improve the efficacy of antimicrobial therapy for difficult-to-treat infections, including overcoming concerns of antimicrobial resistance.<sup>11-13</sup>

Recent studies have shown that using fungal calcineurin pathways holds great promise for the future development of novel agents, including combination therapy with antifungals against fungal pathogens.<sup>14-17</sup> The antifungal activity obtained by the combination of the calcineurin inhibitor tacrolimus (FK506) plus antifungal agents has not yet been evaluated against *Trichosporon* species. In this context, the aim of this study was to evaluate the *in vitro* activity of the combination of FK506 with amphotericin B, fluconazole, itraconazole, and caspofungin against fluconazole-susceptible and fluconazole-resistant *T. asahii* strains.

## Material and methods

### Clinical isolates and molecular identification

One group of 30 fluconazole-susceptible (FS) strains of clinical isolates *T. asahii* maintained in the collection of the Department of Microbiology and Parasitology at the Federal University of Santa Maria, Santa Maria, RS, Brazil were studied. A second group of fluconazole-resistant strains (FR) ( $n=30$ ) was obtained from the FS group after sequential exposure to growing concentrations of fluconazole, as previously described by Fekete-Forgacs et al.,<sup>18</sup> with the following modifications: the final concentration of fluconazole was  $128\mu\text{g mL}^{-1}$ , and the incubation temperature was  $35^\circ\text{C}$  with shaking for 48 h. Cells from this culture were plated on SDA plates, and a single colony was designated isolated FR. The standard strain *T. asahii* CBS 2479 was also included in the susceptibility tests.

The identity of these isolates was confirmed using standard microbiological and molecular methods. Total DNA was extracted according to the protocol described by Moller et al.<sup>19</sup> and Klassen et al.<sup>20</sup> with modifications. Amplification of the IGS1 region (rDNA intergenic) was performed by PCR using the primers 26F (5'ATCCTTTGCAGACGACTTGA-3') and 5SR (5'AGCTTGACTTCGCAGATCGG-3').<sup>5</sup> The PCR products were purified and sequenced. These sequences have been deposited in GenBank (<http://www.ncbi.nlm.nih.gov/genbank/>) with the

following accession numbers: KR233249, KT438844, KR872655, KR872656, KR872657, KR872660, KR872661, KR233246, KR233247, KR233248, KT365854, KT365855, KR872662, KT438839, KT438840, KR872663, KR912056, KT438842, KT365856, KT438847, KR912058, KR912064, KT365858, KR912065, KT365859, KT438846, KT365860, KR912059, KT365861, KR912066.

### Chemicals

Amphotericin B (AMB) (Sigma Chemical Co. – St. Louis, MO, USA), fluconazole (FCZ) (Sigma Chemical Co. – St. Louis, MO, USA), itraconazole (ITZ) (Frangon of Brazil, Pharmaceutical, Ltd., São Paulo, Brazil), caspofungin (CAS) (Merck, Darmstadt, Germany), and tacrolimus (FK506) (Janssen-Cilag Pharmaceutica, Beerse, Belgium) were employed.

The stock solutions of the drugs were prepared in dimethyl sulfoxide (DMSO, Sigma Chemical Co.), except for FCZ, which was diluted in sterile distilled water. FK506 was dissolved in methanol. Working solutions were prepared according to the document M27-A3 of the Clinical and Laboratory Standards Institute.<sup>21</sup>

### *In vitro* susceptibility and drug interaction tests

Susceptibility assays were performed using the broth microdilution method, as described by the document M27-A3 of the Clinical and Laboratory Standards Institute.<sup>21</sup> The strains *Candida krusei* ATCC 6258 and *Candida parapsilosis* ATCC 22019 were included in the tests as controls. The minimum inhibitory concentrations (MICs) were defined as the lowest drug concentration able to inhibit 50% (for FCZ, ITZ, and CAS) or 100% (for AMB and FK506) of fungal growth when compared to the growth of the control. The highest concentrations used were  $16\mu\text{g mL}^{-1}$  for AMB,  $128\mu\text{g mL}^{-1}$  for FCZ,  $16\mu\text{g mL}^{-1}$  for ITZ and  $64\mu\text{g mL}^{-1}$  for CAS and FK506. The MICs employed as indicators for resistance were:  $\geq 2\mu\text{g mL}^{-1}$  (AMB)<sup>5,22,23</sup>;  $\geq 64\mu\text{g mL}^{-1}$  (FCZ)<sup>6,24</sup>;  $\geq 1\mu\text{g mL}^{-1}$  (ITZ),<sup>23</sup> and  $\geq 2\mu\text{g mL}^{-1}$  (CAS).<sup>5,23</sup> After exposing the strains to growing concentrations of FLZ, the strains were separated in two groups: (a) FS group (fluconazole-susceptible group), formed by the strains not exposed to FLZ; and (b) FR group (fluconazole-resistant group) formed by the same strains after exposure to FLZ, and now showing MICs ranging from 64 to  $128\mu\text{g mL}^{-1}$ .

*In vitro* combinations of antifungal agents with FK506 against the FS and FR groups of *T. asahii* were evaluated by the microdilution checkerboard method.<sup>25</sup> For the calculation of the fractional inhibitory concentration index (FICI), MIC values related to 100% inhibition of growth were used. Synergism was defined as the  $\text{FICI} \leq 0.5$ , indifference was  $0.5 < \text{FICI} \leq 4.0$ , and antagonism  $\text{FICI} > 4.0$ . The FICIs were calculated for all wells along the turbidity/non-turbidity interface where the lowest FICI as the final point was determined.

### Statistical analysis

The statistical analysis used to test the Susceptibility of the two groups (susceptible strains vs. resistant strains) to treatment with individual antifungal agents was analyzed with the t-test. Statistical significance was set at  $p < 0.05$ . Results

were analyzed using the software Graph Pad Prism5 (Graph Pad Software version 6.01, CA).

## Results

The *in vitro* susceptibilities of 30 *T. asahii* isolates (FS) against the antifungal agents and FK506 are described in Table 1. FS strains showed low MICs range for FCZ ( $1.0\text{--}16.0\ \mu\text{g mL}^{-1}$ ) and ITZ ( $0.13\text{--}1.0\ \mu\text{g mL}^{-1}$ ) compared to FR strains: FCZ ( $64.0$  to  $>128.0\ \mu\text{g mL}^{-1}$ ) and ITZ ( $0.5\text{--}16.0\ \mu\text{g mL}^{-1}$ ). Interestingly, for AMB, the number of resistant isolates decreased (90–3.33%) in the FR group (MIC range =  $0.13\text{--}4.0\ \mu\text{g mL}^{-1}$ ); and for CAS the MICs remained similar in the two groups (MIC range for FS or FR =  $4.0\text{--}16.0\ \mu\text{g mL}^{-1}$ ). FK506 did not show antifungal activity against FS and FR isolates at the highest concentration tested (MICs  $>64.0\ \mu\text{g mL}^{-1}$ ). The statistical analysis showed significant differences between the susceptibility of FS vs. FR groups for azole antifungal agents (FLZ:  $p < 0.0001$ , ITZ:  $p < 0.007$ ), as well as for AMB ( $p < 0.0001$ ) and CAS ( $p < 0.0001$ ).

The MICs and combinations results of antifungal agents and tacrolimus against *T. asahii* before and after resistance induction are presented in Tables 2 and 3. FK506 combined with AMB against the FS *T. asahii* isolates showed the highest percentage of synergism (96.67%), followed by the combination with CAS (73.33%). In the synergistic interactions between AMB + FK506 or CAS + FK506, the MIC values of FK506 decreased from  $64\ \mu\text{g mL}^{-1}$  to  $0.5\ \mu\text{g mL}^{-1}$ . In contrast, in the FR group the majority of combinations with FK506 showed interactions classified as indifferent: AMB (76.67%), CAS (73.33%), FCZ (63.33%) and ITZ (50%). Antagonisms were not detected for combinations with FCZ and CAS but were showed with AMB (13.33%) and ITZ (10%).

## Discussion

*T. asahii* is the most frequently involved species in disseminated fungal infections including cases of systemic infection in transplanted patients related to therapeutic failures.<sup>2,3</sup> The prognosis of trichosporonosis is very poor, showing mortality rates as high as 80%.<sup>1,10</sup> The lack of response to therapy is related to the relative resistance of *T. asahii* to many different antifungals.<sup>3,4,7,9</sup>

Moreover, the correct identification of *Trichosporon* species is important because the susceptibility is species dependent. In agreement with previous reports,<sup>10,26,27</sup> our results confirm that *T. asahii* clinical isolates seem to be more resistant *in vitro* to amphotericin B than to triazole compounds (Table 1).

Interestingly, our findings demonstrated that after sequential exposure to growing concentrations of fluconazole, the number of resistant isolates to AMB (MIC values  $\geq 2\ \mu\text{g mL}^{-1}$ ) decreased from 90% to 3.33% (Table 1). Although this inverse relationship has been previously reported,<sup>5,7,27,28</sup> targeted deletion of *ERG3* or *ERG11* in *Candida albicans* and *Candida glabrata* seems to be a potential mechanism that may lead to increased susceptibility to AMB with increased resistance to azoles.<sup>29,30</sup>

Similarly, clinical failure and breakthrough infections with *Trichosporon* have been reported with the use of echinocandins.<sup>3,31–34</sup> As expected, we found MICs  $\geq 4\ \mu\text{g mL}^{-1}$  for caspofungin in all our isolates (Table 1). The cases of intrinsic resistance to echinocandins already described for *Cryptococcus* spp., *Trichosporon* spp., *Fusarium* spp., and zygomycetes are associated with insufficient sensitivity of the target enzyme, beta 1,3-D-glucan synthase, to the drug or a mutated form of the enzyme that precludes echinocandin binding.<sup>35</sup>

Regarding the azole antifungal agents, our results demonstrated that prolonged exposure of the clinical isolates to fluconazole showed an increased MIC for itraconazole characterizing cross-resistance among azoles (Table 1). Multidrug resistance to antifungal agents has been reported for *T. asahii* in previous studies.<sup>4,7,9</sup> *T. asahii* clinical isolates from nongranulocytopenic patients showed reduced susceptibility *in vitro* to AMB, flucytosine, fluconazole, itraconazole and ketoconazole.<sup>7</sup> Kushima et al.<sup>9</sup> described that long term use of fluconazole *in vivo* may lead to replacement of the amino acid *ERG11p* and thus trigger *T. asahii* resistance to multiple drugs. This study demonstrated that the MICs for other azole antifungal agents against *T. asahii* strains increased in parallel with the MIC for fluconazole, while the AMB MICs did not significantly change.<sup>9</sup>

Tacrolimus (previously known as FK506) is an effective immunosuppressant, obtained from *Streptomyces tsukubaensis*, and is widely used for prevention of transplant rejection.<sup>36</sup>

**Table 1 – *In vitro* susceptibility of *Trichosporon asahii* isolates to antifungal agents and tacrolimus.**

Agents	Group of isolates	Geometric mean	MIC range	MIC <sub>50</sub>	MIC <sub>90</sub>	Number (%) of resistant isolates
AMB	FS	2.24	1.0–8.0	2.0	4.0	27 (90)
	FR	0.41	0.13–4.0	0.25	1.0	1 (3.33)
FCZ	FS	2.41	1.0–16.0	2.0	8.0	0 (0)
	FR	64.00	64.0 to $>128.0$	64.0	64.0	30 (100)
ITZ	FS	0.27	0.13–1.0	0.25	0.50	1 (3.33)
	FR	1.52	0.5–16.0	1.0	4.0	27 (90)
CAS	FS	7.82	4.0–16.0	8.0	8.0	30 (100)
	FR	7.29	4.0–16.0	8.0	8.0	30 (100)
FK506	FS	ND	$>64.0$	ND	ND	ND
	FR	ND	$>64.0$	ND	ND	ND

MIC range, minimal inhibitory concentration range ( $\mu\text{g mL}^{-1}$ ); MIC<sub>50</sub>, MIC at which 50% of isolates tested were inhibited; MIC<sub>90</sub>, MIC at which 90% of isolates tested were inhibited; AMB, amphotericin B; FCZ, fluconazole; ITZ, itraconazole; CAS, caspofungin; FK506, tacrolimus; FS e FR, fluconazole-susceptible and fluconazole-resistant *Trichosporon asahii* isolates; ND, not determined.

**Table 2 – Minimal inhibitory concentrations (MICs) and combinations results of antifungal agents and tacrolimus against *Trichosporon asahii* before resistance induction.**

Isolates	MICs ( $\mu\text{g mL}^{-1}$ ) and combinations results before resistance induction										
	AMB/FK506			FCZ/FK506			ITZ/FK506			CAS/FK506	
	On its own	In the combination		On its own	In the combination		On its own	In the combination		On its own	In the combination
06	2.00	0.50	S	8.00	1.00	I	1.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
11	2.00	0.25	S	2.00	1.00	I	1.00	2.00	I	16.00	8.00 I
	>64.00	0.50		>64.00	8.00		>64.00	0.50		>64.00	0.50
13	2.00	0.50	S	4.00	1.00	S	1.00	1.00	I	16.00	8.00 I
	>64.00	0.50		>64.00	4.00		>64.00	0.50		>64.00	0.50
19	2.00	0.25	S	4.00	1.00	S	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	16.00		>64.00	0.50		>64.00	0.50
21	2.00	0.50	S	32.00	1.00	I	2.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	64.00		>64.00	0.50		>64.00	0.50
29	4.00	1.00	S	64.00	1.00	I	2.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	64.00		>64.00	0.50		>64.00	0.50
31	2.00	0.25	S	4.00	1.00	S	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	16.00		>64.00	0.5		>64.00	0.50
32	4.00	1.00	S	16.00	1.00	I	2.00	1.00	I	32.00	16.00 I
	>64.00	0.50		>64.00	64.00		>64.00	0.50		>64.00	0.50
36	1.00	0.13	S	8.00	1.00	I	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
43	2.00	0.25	S	64.00	1.00	S	1.00	0.25	S	32.00	4.00 S
	>64.00	0.50		>64.00	4.00		>64.00	0.50		>64.00	0.50
44	4.00	0.25	S	8.00	1.00	I	1.00	2.00	I	32.00	16.00 I
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
46	2.00	0.50	S	4.00	1.00	I	1.00	1.00	I	32.00	16.00 I
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
47	8.00	0.50	S	32.00	1.00	I	2.00	1.00	I	64.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
48	2.00	0.50	S	8.00	1.00	I	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
49	2.00	0.25	S	16.00	1.00	S	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	16.00		>64.00	0.50		>64.00	0.50
50	2.00	0.25	S	8.00	1.00	S	1.00	1.00	I	32.00	16.00 I
	>64.00	0.50		>64.00	16.00		>64.00	0.50		>64.00	0.50
51	4.00	0.25	S	2.00	1.00	I	1.00	1.00	I	32.00	16.00 I
	>64.00	0.50		>64.00	8.00		>64.00	0.50		>64.00	0.50
53	4.00	0.25	S	4.00	1.00	S	2.00	0.50	S	16.00	16.00 I
	>64.00	0.50		>64.00	16.00		>64.00	0.50		>64.00	0.50
54	2.00	0.50	S	8.00	1.00	I	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
56	2.00	0.50	S	8.00	1.00	I	1.00	0.50	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
58	2.00	0.50	S	16.00	1.00	I	1.00	1.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
59	1.00	0.50	I	16.00	1.00	S	1.00	0.25	S	64.00	8.00 S
	>64.00	0.50		>64.00	8.00		>64.00	0.50		>64.00	0.50
60	2.00	0.25	S	8.00	1.00	S	1.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	16.00		>64.00	0.50		>64.00	0.50
61	8.00	0.50	S	8.00	1.00	I	1.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
62	2.00	0.25	S	8.00	1.00	I	1.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
63	2.00	0.50	S	8.00	1.00	S	1.00	0.02	I	32.00	8.00 S
	>64.00	0.50		>64.00	16.00		>64.00	64.00		>64.00	0.50
64	2.00	0.25	S	4.00	1.00	S	0.50	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	8.00		>64.00	0.50		>64.00	0.50
65	4.00	0.25	S	8.00	1.00	I	0.50	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	32.00		>64.00	0.50		>64.00	0.50
66	2.00	0.50	S	8.00	1.00	I	1.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	64.00		>64.00	0.50		>64.00	0.50
67	1.00	0.25	S	4.00	1.00	S	1.00	2.00	I	32.00	8.00 S
	>64.00	0.50		>64.00	16.00		>64.00	0.50		>64.00	0.50

MICs, minimal inhibitory concentrations as the lowest concentration that showed 100% inhibition of fungal growth ( $\mu\text{g mL}^{-1}$ ); AMB, amphotericin B; FCZ, fluconazole; ITZ, itraconazole; CAS, caspofungin; FK506, tacrolimus; S, synergism; I, indifference.

**Table 3 – Minimal inhibitory concentrations (MICs) and combinations results of antifungal agents and tacrolimus against *Trichosporon asahii* after resistance induction.**

Isolates	MICs ( $\mu\text{g mL}^{-1}$ ) and combinations results after resistance induction											
	AMB/FK506			FCZ/FK506			ITZ/FK506			CAS/FK506		
	On its own	In the combination		On its own	In the combination		On its own	In the combination		On its own	In the combination	
06	1.00	0.50	I	>128.00	32.00	S	>16.00	2.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
11	0.25	0.25	I	128.00	32.00	S	4.00	1.00	S	8.00	4.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
13	0.50	0.50	I	>128.00	64.00	I	>16.00	4.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
19	0.50	0.25	I	128.00	64.00	I	4.00	2.00	I	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
21	0.25	0.25	I	128.00	32.00	S	2.00	1.00	I	32.00	8.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
29	1.00	0.25	S	128.00	128.00	I	>16.00	8.00	I	16.00	32.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
31	0.50	0.13	S	128.00	128.00	I	4.00	0.50	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
32	0.25	0.50	I	128.00	128.00	I	>16.00	4.00	S	16.00	32.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
36	0.50	0.25	I	128.00	64.00	I	2.00	2.00	I	16.00	4.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
43	0.25	0.25	I	128.00	32.00	S	2.00	1.00	I	16.00	4.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
44	0.25	0.25	I	128.00	32.00	S	>16.00	2.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
46	1.00	1.00	I	128.00	128.00	I	>16.00	2.00	S	16.00	16.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
47	4.00	0.25	S	64.00	64.00	I	2.00	2.00	I	16.00	16.00	I
	>64.00	5.00		>64.00	0.50		>64.00	0.50		>64.00	0.50	
48	0.50	0.25	I	>128.00	32.00	S	4.00	1.00	S	16.00	4.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
49	0.50	0.25	I	128.00	64.00	I	8.00	2.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
50	0.50	0.25	I	128.00	32.00	S	>16.00	1.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
51	0.25	0.25	I	128.00	32.00	S	2.00	2.00	I	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
53	0.25	0.50	I	128.00	128.00	I	2.00	16.00	A	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
54	0.25	0.13	I	128.00	64.00	I	>16.00	1.00	I	16.00	16.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
56	0.13	0.50	A	>128.00	64.00	I	>16.00	8.00	I	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
58	0.50	2.00	A	128.00	128.00	I	8.00	32.00	A	16.00	1.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
59	0.50	0.25	I	>128.00	64.00	I	4.00	2.00	I	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
60	0.25	0.50	I	128.00	128.00	I	>16.00	32.00	I	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
61	0.25	0.50	I	>128.00	64.00	I	>16.00	4.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
62	1.00	0.50	I	>128.00	128.00	I	4.00	2.00	I	16.00	16.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
63	0.25	2.00	A	>128.00	8.00	S	16.00	8.00	I	16.00	2.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
64	0.25	0.13	I	128.00	64.00	I	2.00	2.00	I	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
65	0.25	1.00	A	>128.00	16.00	S	1.00	8.00	A	16.00	1.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
66	0.25	0.50	I	128.00	64.00	I	1.00	2.00	I	16.00	2.00	S
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	
67	0.50	0.50	I	128.00	32.00	S	8.00	2.00	S	16.00	8.00	I
	>64.00	0.50		>64.00	0.50		>64.00	0.50		>64.00	0.50	

MICs, minimal inhibitory concentrations as the lowest concentration that showed 100% inhibition of fungal growth ( $\mu\text{g mL}^{-1}$ ); AMB, amphotericin B; FCZ, fluconazole; ITZ, itraconazole; CAS, caspofungin; FK506, tacrolimus; S, synergism; I, indifference; A, antagonism.



This compound exerts its effects by blocking the immune system through inhibition of calcineurin.<sup>37</sup> Moreover, this protein can also affect essential functions of the fungal cell and it is intrinsically involved in the growth and pathogenesis of three major fungal species: *Cryptococcus neoformans*, *C. albicans*, and *Aspergillus fumigatus*.<sup>36,38</sup> In this study, FK506 showed low antifungal activity when tested alone against FS and FR *T. asahii* isolates (MICs > 64.0 µg mL<sup>-1</sup>).

However, against the FS group, the results of our study (Table 2) demonstrate strong in vitro synergism of FK506 combined with drugs that were ineffective in inhibiting this group of *T. asahii* clinical isolates (Table 1) such as caspofungin (73.33%) and AMB (96.67%). High percentage synergistic interactions for caspofungin plus FK506 have been previously reported against *C. neoformans*, *Fusarium* spp., and *Aspergillus* spp.<sup>11,14,39</sup>

The potential enhanced antifungal activity of caspofungin in combination with other antifungal agents and anti-calcineurin drugs against clinical isolates of *Fusarium* spp. was demonstrated by Shalit et al.<sup>11</sup> The association of this echinocandin with FK506 appeared synergistic against all the isolates tested.<sup>11</sup> The antifungal effect exhibited by immunosuppressants cyclosporin A and FK506 is probably related to calmodulin activated protein phosphatase involved in fungal stress response, virulence, and antifungal resistance.<sup>36</sup> Steinbach et al.<sup>14</sup> also demonstrated a synergistic interaction of cyclosporin and FK506 with caspofungin against *A. fumigatus*. In this study, the calcineurin inhibitors were capable of causing a delay in filamentation of *A. fumigatus*, which suggested that inhibition of this pathway may potentiate the action of conventional antifungal agents in combination therapy against invasive aspergillosis.<sup>14</sup>

The pharmacokinetics of caspofungin is unaltered by coadministration of tacrolimus, but caspofungin may reduce tacrolimus concentrations by up to 20%.<sup>40</sup> Therefore, monitoring standard tacrolimus blood concentrations and appropriate tacrolimus dose adjustments are recommended for patients receiving both therapies.

On the other hand, the synergism observed with the association of FK506 and AMB can benefit the antifungal therapy regimen through the reductions in time to treatment response, dose with associated toxicity, costs, and decreased potential of microorganism-acquired resistance. Although AMB has been reported to have an insignificant effect on tacrolimus metabolism,<sup>41</sup> this polyene is well known to be nephrotoxic. Therefore, monitoring of serum creatinine is probably warranted for patients receiving both drugs.<sup>42</sup>

Studies have also shown that calcineurin inhibitors may exert a synergistic interaction when combined with antifungal azoles against *Candida* species.<sup>15,16</sup> However, our results demonstrated that combinations of FCZ or ITZ with FK506 produced interactions mainly indifferent emphasizing a lack of effect against FS and FR *T. asahii* isolates (Tables 2 and 3). The pharmacokinetics interactions between FK506 and azoles are well known and show that azoles inhibit the metabolism of FK506, requiring monitoring of the plasmatic concentration of FK506.<sup>42</sup>

In conclusion, our findings demonstrated that the combination of FK506 with AMB or CAS leads to high rates of synergism in vitro. These combinations against *T. asahii*

isolates that presented resistance to both AMB and CAS deserve attention as candidates for in vivo studies focusing *T. asahii* experimental infections.

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgments

Alves SH thanks the financial support provided by the Brazilian Agencies FAPERGS (Grant Proc. 2261-12) and CNPq (Grant Proc. 470229/2012-8).

## REFERENCES

- Girmeria C, Pagano L, Martino B, et al. Invasive infections caused by *Trichosporon* species and *Geotrichum capitatum* in patients with hematological malignancies: a retrospective multicenter study from Italy and review of the literature. *J Clin Microbiol.* 2005;43:1818-28.
- Biasoli MS, Carlson D, Chiganeri GJ. Systemic infection caused by *Trichosporon asahii* in a patient with liver transplant. *Med Mycol.* 2008;46:719-23.
- Yang MF, Gao HN, Li LJ. A fatal case of *Trichosporon asahii* fungemia and pneumonia in a kidney transplant recipient during caspofungin treatment. *Ther Clin Risk Manag.* 2014;10:759-62.
- Falk R, Wolf DG, Shapiro M, Polacheck I. Multidrug-resistant *Trichosporon asahii* isolates are susceptible to voriconazole. *J Clin Microbiol.* 2003;41:911.
- Chagas-Neto TC, Chaves GM, Melo ASA, et al. Bloodstream infections due to *Trichosporon* spp.: species distribution. *Trichosporon asahii* genotypes determined on the basis of ribosomal DNA intergenic spacer 1 sequencing, and antifungal susceptibility testing. *J Clin Microbiol.* 2009;47:1074-81.
- Harizolan G, Canton E, Sahin S, et al. Head-to-head comparison of inhibitory and fungicidal activities of fluconazole, itraconazole, voriconazole, posaconazole, and isavuconazole against clinical isolates of *Trichosporon asahii*. *Antimicrob Agents Chemother.* 2013;57:4841-7.
- Wolf DG, Falk R, Hacham M, et al. Multidrug-resistant *Trichosporon asahii* infection of nongranulocytopenic patients in three intensive care units. *J Clin Microbiol.* 2001;39:4420-5.
- Pfaller MA, Diekema DJ, Gibbs DL, et al. Results from the ARTEMIS DISK Global Antifungal Surveillance Study, 1997 to 2007: 10.5-year analysis of susceptibilities of noncandidal yeast species to fluconazole and voriconazole determined by CLSI standardized disk diffusion testing. *J Clin Microbiol.* 2009;47:117-23.
- Kushima H, Tokimatsu I, Ishii H, et al. Cloning of the lanosterol 14- $\alpha$ -demethylase (ERG11) gene in *Trichosporon asahii*: a possible association between G453R amino acid substitution and azole resistance in *T. asahii*. *FEMS Yeast Res.* 2012;12:662-7.
- Colombo AL, Padovan ACB, Chaves GM. Current knowledge of *Trichosporon* spp. and trichosporonosis. *Clin Microbiol Rev.* 2011;24:682-700.
- Shalit I, Shadkchan Y, Mercus G. In vitro synergy of caspofungin with licensed and novel antifungal drugs against clinical isolates of *Fusarium* spp. *Med Mycol.* 2009;47:457-62.

12. Venturini TP, Chassot F, Loreto ES, et al. Antifungal activities of diphenyl diselenide and ebselen alone and in combination with antifungal agents against *Fusarium* spp. *Med Mycol*. 2016;1-6.
13. Scheid LA, Mario DAN, Kubiça TF, et al. *In vitro* activities of antifungal agents alone and in combination against fluconazole-susceptible and -resistant strains of *Candida dubliniensis*. *Braz J Infect Dis*. 2012;16:78-81.
14. Steinbach WJ, Schell WA, Blakeship JR, et al. *In vitro* interactions between antifungals and immunosuppressants against *Aspergillus fumigatus*. *Antimicrob Agents Chemother*. 2004;48:1664-9.
15. Sun S, Li Y, Guo Q, et al. *In vitro* interactions between tacrolimus and azoles against *Candida albicans* determined by different methods. *Antimicrob Agents Chemother*. 2008;52:409-17.
16. Uppuluri P, Heitman J, Andes D, et al. Synergistic effect of calcineurin inhibitors and fluconazole against *Candida albicans* biofilms. *Antimicrob Agents Chemother*. 2008;52:1127-32.
17. Denardi LB, Mario DAN, Loreto ES, et al. Synergistic effects of tacrolimus and azole antifungal compounds in fluconazole-susceptible and fluconazole-resistant *Candida glabrata* isolates. *Braz J Microbiol*. 2015;46:125-9.
18. Fekete-Forgács K, Gyurc L, Lenkey B. Changes of virulence factors accompanying the phenomenon of induced fluconazole resistance in *Candida albicans*. *Mycoses*. 2000;43:273-9.
19. Moller EM, Bahnweg G, Sandemann H, et al. A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Res*. 1992;20:6115-6.
20. Klassen GR, Balcerzak M, De Cock AWAM. 5S ribosomal RNA gene spacer as species specific probes for eight species of *Pythium*. *Phytopathology*. 1996;86:581-7.
21. Clinical and Laboratory Standards Institute. Reference method for broth dilution antifungal susceptibility testing of yeasts. Approved standard M27-A3. Wayne, USA: CLSI; 2008.
22. Montoya AM, González AS, Palma-Nicolás JP, et al. Genotyping, extracellular compounds, and antifungal susceptibility testing of *Trichosporon asahii* isolated from Mexican patients. *Med Mycol*. 2015;53:505-11.
23. Ribeiro MA, Alastruey-Izquierdo A, Gomez-Lopez A, et al. Molecular identification and susceptibility testing of *Trichosporon* isolates from a Brazilian hospital. *Rev Iberoam Micol*. 2008;25:221-5.
24. Menezes EA, Marinho JAS, Angelo MRF, et al. Isolation and antifungal Susceptibility testing of *Trichosporon asahii* in Ceará, Brazil. *Rev Inst Med Trop Sao Paulo*. 2012;54:1-3.
25. Moody J. Synergism testing: broth microdilution checkerboard and broth macrodilution methods. In: Garcia LS, Isenberg HD, editors. *Clinical microbiology procedures handbook*. 2nd ed. Washington, DC: ASM Press; 2007.
26. Paphitou NI, Ostrosky-Zeichner L, Paetznick VL, et al. *In vitro* antifungal susceptibilities of *Trichosporon* species. *Antimicrob Agents Chemother*. 2002;46:1144-6.
27. Rodriguez-Tudela JL, Diaz-Guerra TM, Mellado E, et al. Susceptibility patterns and molecular identification of *Trichosporon* species. *Antimicrob Agents Chemother*. 2005;49:4026-34.
28. Kalkanci A, Sugita T, Arkan S, et al. Molecular identification, genotyping, and drug susceptibility of the basidiomycetous yeast pathogen *Trichosporon* isolated from Turkish patients. *Med Mycol*. 2010;48:141-6.
29. Sanglard D, Ischer F, Parkinson T, et al. *Candida albicans* mutations in the ergosterol biosynthetic pathway and resistance to several antifungal agents. *Antimicrob Agents Chemother*. 2003;47:2404-12.
30. Cho EJ, Shin JH, Kim SH. Emergence of multiple resistance profiles involving azoles, echinocandins and amphotericin B in *Candida glabrata* isolates from a neutropenia patient with prolonged fungaemia. *J Antimicrob Chemother*. 2014;70:1268-70.
31. Bayramoglu G, Sonmez M, Tosun I, et al. Breakthrough *Trichosporon asahii* fungemia in neutropenic patient with acute leukemia while receiving caspofungin. *Infection*. 2008;36:68-70.
32. Karapinar DY, Karadas N, Sivas ZO, et al. Rare severe mycotic infections in children receiving empirical caspofungin treatment for febrile neutropenia. *Braz J Infect Dis*. 2015;19:549-52.
33. Liao Y, Harmann T, Zheng T, Yang RY, Ao JH, Wang WL. Breakthrough trichosporonosis in patients receiving echinocandins: case report and literature review. *Chin Med J*. 2012;125:2632-5.
34. Sun HY, Singh N. Characterization of breakthrough invasive mycoses in echinocandin recipients: an evidence-based review. *Int J Antimicrob Agents*. 2010;35:211-8.
35. Chandra J, Mohammed S, Ghannoum MA. Echinocandins: exploring susceptibility and resistance. In: Mayers DL, editor. *Antimicrobial drug resistance*. New York: Humana Press; 2009.
36. Steinbach WJ, Reedy JL, Cramer RA Jr, et al. Harnessing calcineurin as a novel anti-infective agent against invasive fungal infections. *Nat Rev Microbiol*. 2007;5:418-30.
37. Baksh S, Burakoff SJ. The role of calcineurin in lymphocyte activation. *Semin Immunol*. 2000;12:405-15.
38. Blankenship JR, Heitman J. Calcineurin is required for *Candida albicans* to survive calcium stress in serum. *Infect Immun*. 2005;73:5767-74.
39. Kontoyiannis DP, Lewis RE, Oshero N, et al. Combination of caspofungin with inhibitors of the calcineurin pathway attenuates growth *in vitro* in *Aspergillus* species. *J Antimicrob Chemother*. 2003;51:313-6.
40. Sable CA, Nguyen BY, Chodakewitz JA, DiNubile MJ. Safety and tolerability of caspofungin acetate in the treatment of fungal infections. *Transplant Infect Dis*. 2002;4:25-30.
41. Iwasaki K, Matsuda H, Nagase K, et al. Effects of twenty-three drugs on the metabolism of FK506 by human liver microsomes. *Res Comm Chem Pathol Pharmacol*. 1993;82:209-16.
42. Paterson DL, Singh N. Interactions between tacrolimus and antimicrobial agents. *Clin Infect Dis*. 1997;25:1430-40.